

1/6




FIGURE 1

BEST AVAILABLE COPY

2/6

A. The amino terminal sequence of IL13-gC

gcttgggtcgggagggccgcacatcgaacgcacaccccccatccgggtgggtccgtgtggaggtcggtttttcagtgcc
 cggctctcgctttgcccgggaacgctagcctcATGGCGCTTTTGTGTGACCACGGTCATTGCTCTCACTTGCC
 gC upstream. IL-13*
 TGGCGGCTTTGCCTCCCCAGGCCCTGTGCCCTCCCTCTACAGCCCTCAGGTAGCTCATTGAGGAGCTGGTCA
 ACATCAGCCAGAACCCAGAAGGCTCCGCTCTGCAATGGCAGCATGGTATGGAGCATCAACCTGACAGCTGGC
 ATGTACTGTGCAGCCCTGGAATCCCTGATCAACGTGTCAAGGCTGCAGTGGCATCGAGAAGACCCAGAGGAT
 GCTGAGCGGATTCTGCCCGCACAAAGGTCTCAGCTGGGCAGTTTTTCAGCTTGCAATGTCCGAGACACCAAAA
 TCGAGGTGGCCAGTTTGTAAAAGATCTGCTCTTACATTTAAAGAACTTTTTTCGCGAGGGACGGTTgaa
 tgcaccccgcatggagttccgcctccagatatggcggttactccatgggtccgtcccccccaatcgctccggc
 gC downstream

B. The sequence of the gB_{Δpoly(K)} domain


GGGTCTGGTGGCGTCGGCGGCTCCGAGTTCCCCGGCACGCCTGGGGTCGGGGCCGCGACCCAGGCGGC
 GAACGGGGGACCTGCCACTCCGGCGCCGCCCCGCCCTGGCCCCGCCCCAACGGGGGATCCGAAACCGAAG
 AGAAGCAGAAAACCGAAACCCCAAGCGCCCGCGCCCCGCGGGGACAAACCGACCGTCGCCCGGGGCCA
 CGCCACCTTCGCGGAGCACCTGCGGGACATCAAGGCGGAGAACACCGATGCAAACTTTTACGTGTGCCCA
 CCCCCACGGGCGCCACGGTGGTGCAGTTCGAGCAGCCGCGCCGCTGCCCCGACCCGCGCCCGAGGGTCA

C. The amino terminal sequence of IL13-gD

ATGGGGGGGGCTGCCGCCAGGTTGGGGGCGGTGATTTTGTGTGTCGTATAGTCGGCCTC
 Signal peptide of gD→
CATGGGGTCCGCGGCAAATATGCCTTGGCGGATGCCTCTCTCAAGCTGGCCGACCCCAAT
 ←
 CGCTTTCGCCGCAAAGACCTTCCGGTCCctegag*ATGGCGCTTTTGTGTGACCACGGTCATT
24AA XhoI IL13→
 GCTCTCACTTGCCCTTGGCGGCTTTGCCTCCCCAGGCCCTGTGCCTCCCTCTACAGCCCTC
 AGGGAGCTCATTGAGGAGCTGGTCAACATCAGCCAGAACCCAGAAGGCTCCGCTCTGCAAT
 GGCAGCATGGTTTGGAGCATCAACCTGACAGCTGGCATGTACTGTGCAGCCCTGGAATCC
 CTGATCAACGTGTGAGGCTGCAGTGCCATCGAGAAGACCCAGAGGATGCTGGGCGGATTC
 TGGCCGCACAAGGTCTCAGCTGGGCAAGTTTTCCAGCTTGCAATGTCCGAGACACCAAAATC
 GAGGTGGCCAGTTTGTAAAGGACCTGCTCTTACATTTAAAGAACTTTTTTCGCGAGGGA
 CGGTTCAACTGAAAC*ggtaaccCTGGACCAGCTGACCGACCCCTCCGGGGGTCCGGCGCGTG
←IL13 KpnI 25AA
 TACCACATCCAGGCGGGCCTACCGGACCCGTTCCAGCCCCCAGCCTCCCGATC

FIGURE 2

BEST AVAILABLE COPY

**FIGURE 3**

BEST AVAILABLE COPY

4/6




FIGURE 4

BEST AVAILABLE COPY

5/6





FIGURE 5

BEST AVAILABLE COPY



pgD- in pcDNA3.1(-) with CMV promoter collapsed by NruI/NheI digestion. The fragment containing gD upstream at NotI/BamHI, and gD downstream at XhoI/KpnI in pBluescript II SK was excised by NotI/KpnI and ligated into pcDNA3.1(-) in which the CMV promoter had been collapsed. N, NotI, B, BamHI, P, PstI, E, EcoRI, H, HindIII, C, ClaI, X, XhoI, and K, KpnI.

FIGURE 6